

A guide to refer your patients with prediabetes to an evidence-based diabetes prevention program





Preventing diabetes: Making a difference by linking the clinic with the community.

In the average primary care practice, it's likely that one-third of patients over age 18, and half over age 65, have prediabetes.

Use this guide to help your patients delay or prevent the onset of type 2 diabetes

Prediabetes is a health condition characterized by blood glucose levels that are higher than normal, but not high enough to be diagnosed as diabetes. Prediabetes increases the risk for type 2 diabetes, heart disease and stroke.

Prediabetes is treatable, but only about 10 percent of people who have it are aware that they do. Left untreated up to one-third of people with prediabetes will progress to diabetes within five years.

During that window of time your patients can benefit from a proven lifestyle change intervention that is part of the <u>National Diabetes Prevention Program</u> (National DPP) led by the Centers for Disease Control and Prevention (CDC).

As part of the National DPP, the American Medical Association (AMA) and the CDC are collaborating to create tools and resources that care teams can use to identify patients with prediabetes, and refer eligible patients to in-person or online diabetes prevention programs.

Physicians and care teams from a diverse group of practices helped the AMA and the CDC create the tools in this guide, and have used them in their own practices to:

- Screen and identify patients for prediabetes
- Refer patients to diabetes prevention programs
- Create feedback loops, linking the patient's progress in the diabetes prevention program back to the practice

Part of a national movement

To achieve CDC recognition as part of the National DPP, programs must provide evidence they are following a CDC-approved curriculum and achieving meaningful results with patients. These programs are based on research showing that a year-long, structured lifestyle change intervention reduced the incidence of diabetes by 58 percent among adults with prediabetes and by 71 percent in those aged 60 years or older.

These programs are successful in part because they require only moderate weight loss to achieve preventive health benefits. Weight loss of 5 to 7 percent of body weight—10 to 14 pounds for a person weighing 200 pounds—led to the results mentioned above.

The AMA and the CDC are promoting these diabetes prevention programs because they are one of the most effective ways to help physicians prevent or delay type 2 diabetes in high-risk patients.

Use this implementation guide and its tools to help identify and refer patients with prediabetes to a diabetes prevention program that is part of the CDC's National DPP.







Overview of guide tools

| Resource | Purpose | | | | | | |
|---|---|--|--|--|--|--|--|
| Engage clinicians | | | | | | | |
| You can prevent type 2 diabetes Health care provider fact sheet | Provides a brief overview of the evidence-based diabetes prevention program and a rationale for engaging with the program, such as improved patient outcomes. Also assists clinicians in advocating to their colleagues and leaders about the value of incorporating diabetes prevention screening and referral into their practices. | | | | | | |
| Engage patients | | | | | | | |
| Diabetes Risk Assessments CDC and American Diabetes Association (ADA) questionnaires | Offers an educational opportunity for patients to learn about their risk for prediabetes, and help physicians and care teams identify their patients at great risk. | | | | | | |
| Promoting prediabetes awareness to your patients 8" x 11" poster) | Helps practices increase patient awareness of prediabetes to pave the way for conversations with patients about screening and referral. | | | | | | |
| Are you at risk for type 2 diabetes? Patient handout | For use by physician practices in patient waiting areas to increase patient awareness and pave the way for conversations with patients about screening and referral. | | | | | | |
| So you have prediabetes now what? Patient handout | For use by physician practices in the exam room after screening has revealed that a patient has prediabetes. Helps the patient leave the office visit with concrete information for later reference. | | | | | | |
| Sample "Patient letter/email and phone script" | Enables physician practices to conduct efficient follow-up and referral with patients who have been identified as having prediabetes, informing them of their prediabetes status and referral to an evidence-based diabetes prevention program. | | | | | | |
| Incorporate screening, testing and r | eferral into practice | | | | | | |
| M.A.P. to diabetes prevention for your practice One-page overview | Offers practices a one-page roadmap to applying the elements of the diabetes prevention screening and referral guide. | | | | | | |
| Patient flow process Infographic | Provides a high-level overview of how office staff can facilitate point-of-care identification. | | | | | | |
| Point-of-care prediabetes identification algorithm Infographic and narrative | With a graphic on one side, and narrative on other, the document offers practices an option to adapt/incorporate a prediabetes screening and referral process into their workflow. | | | | | | |
| Retrospective prediabetes identification algorithm Infographic and narrative | With a graphic on one side, and narrative on other, the document offers practices an option to adapt/incorporate an identification and referral process into their electronic health records and generate a registry of patients at risk for type 2 diabetes. | | | | | | |
| Sample patient referral form/table for calculating body mass index | Makes the referral process easier for practices, helps engage the patient (particularly if they sign the optional patient signature box) and prepares diabetes prevention program providers to engage with the patient as well. | | | | | | |
| Commonly used CPT and ICD codes Table | Enables physician practices to obtain reimbursement for prediabetes screening. | | | | | | |
| Connect your clinic with diabetes pr | evention programs | | | | | | |
| Link to sample "Business Associate Agreement" on AMA's website | Provides link to template agreement some practices have used to share information with diabetes prevention program providers. | | | | | | |







Additional information

AMA diabetes prevention initiative

preventdiabetesstat.org

Learn more about the AMA's commitment to preventing type 2 diabetes.

Centers for Disease Control and Prevention's National Diabetes Prevention Program

cdc.gov/diabetes/prevention

Visit this site for detailed information about the CDC's National Diabetes Prevention Program.

National Diabetes Education Program

ndep.nih.gov/am-i-at-risk/

Find educational resources about preventing diabetes for you and your patients.

How does a diabetes prevention program work?

Diabetes prevention programs that are part of the National DPP use lifestyle change interventions that target improving diet, increasing physical activity and achieving moderate weight loss.

The goal for each participant is to lose \geq 5% of body weight by:

- Progressively reducing dietary intake of calories and fat through improved food choices
- Gradually increasing moderate physical activity (e.g., brisk walking) to ≥150 minutes per week
- Developing behavioral problem-solving and coping skills

Features include:

- A year-long structured program (in-person group, online or distance learning) consisting of:
 - An initial six-month phase offering at least 16 sessions over 16–24 weeks
 - A second six-month phase offering at least one session a month (at least six sessions)
- · Facilitation by a trained lifestyle coach
- Use of a CDC-approved curriculum
- · Regular opportunities for direct interaction between the lifestyle coach and participants
- An emphasis on behavior modification, managing stress and peer support

Who is eligible for referral to a diabetes prevention program?

To be eligible for referral, patients must:

- Be at least 18 years old and
- Be overweight (Body Mass Index (BMI) ≥24; ≥22 if Asian) and
- Have a blood test result in the prediabetes range within the past year:
 - Hemoglobin A1C: 5.7-6.4% or
 - Fasting plasma glucose: 100–125 mg/dL or
 - Two-hour plasma glucose (after a 75 gm glucose load): 140–199 mg/dL or
- · Be previously diagnosed with gestational diabetes and
- Have no previous diagnosis of diabetes

Physicians and other health care providers should also use their independent judgment when referring to a diabetes prevention program.







How can patients find a diabetes prevention program near them?

Diabetes prevention programs are available in varied locations such as local YMCAs, wellness centers, faith-based organizations and worksites—as well as in health care facilities. Online versions are also available. Visit nccd.cdc.gov/DDT_DPRP/Registry.aspx to find a program that is part of the CDC's National DPP recognition program.

Does health insurance cover patient participation in a diabetes prevention program?

A growing number of private health insurers offer coverage for patient participation in diabetes prevention programs. Several employers include coverage as part of workplace wellness programs. Costs for a full year of program participation are approximately \$400–\$500. Some program providers offer monthly payment plans and discounts based on ability to pay. The AMA and the CDC continue to advocate for public and private insurance coverage of the diabetes prevention program.

How do I code for prediabetes screening?

Depending on the type of office visit, practices can use several CPT and ICD codes to bill for prediabetes screening and counseling. A list of commonly used <u>CPT and ICD codes</u> is included in this guide.

Feedback from diabetes prevention program to referring clinicians

Most programs send reports of participant progress to referring clinicians after the eighth and 16th group sessions. In addition, participants in the program complete periodic self-evaluations that referring clinicians can request directly from patients.

Sending patient information to a diabetes prevention program provider Business Associate Agreement

Under the U.S. Health Insurance Portability and Accountability Act of 1996 (HIPAA), a HIPAA Business Associate Agreement (BAA) is a contract that protects personal health information in accordance with HIPAA guidelines. Some physician practices may want to explore whether a BAA is needed to exchange information with a diabetes prevention program. (Link to a "Business Associates Agreement" template on AMA's website.)







Engage clinicians















You can prevent type 2 diabetes

Test your patients for prediabetes and refer those at risk to an evidence-based diabetes prevention program

You likely know which of your patients is at high risk for type 2 diabetes. Until now you may not have had a resource to help them stop the progression from prediabetes to diabetes. Now, you do.

The American Medical Association and the Centers for Disease Control and Prevention (CDC) have created a toolkit that can help physician practices screen and refer patients to evidence-based diabetes prevention programs without adding a burden to your practice. Visit **preventdiabetesstat.org** to learn more.

- Progression from prediabetes to diabetes can take as little as five years.
- During that window of time, your patients can benefit from a proven intervention that is part of the CDC's National Diabetes Prevention Program (National DPP).
- Counsel your patients that prediabetes is a potentially reversible condition, and one that you can help them manage effectively by:
 - Screening and Identifying patients for prediabetes
 - Referring them to a program that is part of the CDC's National DPP

This program is evidence-based

- The diabetes prevention program is a lifestyle intervention based on research funded by the National Institutes of Health that showed, among those with prediabetes, a 58 percent reduction in the number of new cases of diabetes overall, and a 71 percent reduction in new cases for those over age 60.
- These results were achieved through reducing calories, increasing physical activity, and a weight loss of just 5 to 7 percent of body weight—10 to 14 pounds for a person weighing 200 pounds.*
- Based on strong evidence of effectiveness in reducing new-onset diabetes, the Community Preventive Services Task Force (thecommunityguide.org) now recommends combined diet and physical activity promotion programs like the National DPP, for people at increased risk of type 2 diabetes.

Program overview

- · The program empowers patients with prediabetes to take charge of their health and well-being.
- Participants meet in groups with a trained lifestyle coach for 16 weekly sessions and 6–8 monthly follow-up sessions.
- These are NOT exercise classes. At these sessions patients learn ways to incorporate healthier eating and moderate physical activity, as well as problem-solving, stress-reduction and coping skills into their daily lives.

See next page to determine which of your patients is eligible for the diabetes prevention program.

In the average primary care practice it's likely one-third of patients over age 18, and half over age 65, have prediabetes.

* Visit http://diabetes.niddk.nih.gov/dm/pubs/preventionprogram to learn more about this research.











Locating a program

- Programs are offered in varied locations such as local YMCAs, community centers, faith-based organizations, hospitals and worksites, and are also available online.
- Find a program for your patients at **cdc.gov/diabetes/prevention**.

Eligibility for the diabetes prevention program

A. Inclusion criteria:

- Current age ≥18 years **and**
- Most recent BMI ≥24 (≥22 if Asian) and
- A positive lab test result within previous 12 months:
 - HbA1C 5.7-6.4% (LOINC code 4548-4) or
 - FPG 100-125 mg/dL (LOINC code 1558-6) or
 - OGTT 140-199 mg/dL (LOINC code 62856-0) or
- History of gestational diabetes (ICD-9: V12.21)

B. Exclusion criteria:

- Current diagnosis of diabetes (ICD-9: 250.xx) or
- Current Insulin use

Consider referring eligible patients:

- At the time of an office visit, and/or
- By generating a list of eligible patients from your electronic health record using a structured query

Physicians and other health care providers should also use their independent judgment when referring to a diabetes prevention program.







Engage patients





Patient risk assessment

ARE YOU AT RISK FOR

TYPE 2 DIABETES? American Diabetes Association.



191+

198+

204+

211+ 218+

225+

232+ 240+

247+ 255+

270+

278+

286+

294+

302+

311+

319+

328+

Weight (lbs.)

143-190

197-261

203-269

209-277

215-285

221-293

227-301

233-310

240-318

246-327

Diabetes Risk Test

1 How old are you?

Less than 40 years (0 points) 40-49 years (1 point) 50-59 years (2 points) 60 years or older (3 points)

Are you a man or a woman?

Man (1 point) Woman (0 points)

If you are a woman, have you ever been diagnosed with gestational diabetes?

> Yes (1 point) No (0 points)

Do you have a mother, father, sister, or brother with diabetes?

> Yes (1 point) No (0 points)

Have you ever been diagnosed with high blood pressure?

> Yes (1 point) No (0 points)

6 Are you physically active? Yes (0 points)

What is your weight status? (see chart at right)

Write your score in the box.



| 4′ 11″ | 124-147 | 148-197 |
|--------|---------|---------|
| 5′ 0″ | 128-152 | 153-203 |
| 5′ 1″ | 132-157 | 158-210 |
| 5′ 2″ | 136-163 | 164-217 |
| 5′ 3″ | 141-168 | 169-224 |
| 5′ 4″ | 145-173 | 174-231 |
| 5′ 5″ | 150-179 | 180-239 |
| 5′ 6″ | 155-185 | 186-246 |
| 5′ 7″ | 159-190 | 191-254 |

164-196

169-202

174-208

179-214

189-226

194-232

200-239

205-245

4' 10" 119-142

5'9"

5' 10"

5' 11"

6' 0"

6' 1"

6' 2"

6' 3"

6' 4"





(1 Point) (2 Points) (3 Points) You weigh less than the amount in the left column (0 points)

Adapted from Bang et al., Ann Intern Med

Original algorithm was validated without

gestational diabetes as part of the model.

If you scored 5 or higher:

You are at increased risk for having type 2 diabetes. However, only your doctor can tell for sure if you do have type 2 diabetes or prediabetes (a condition that precedes type 2 diabetes in which blood glucose levels are higher than normal). Talk to your doctor to see if additional testing is needed.

Type 2 diabetes is more common in African Americans, Hispanics/ Latinos, American Indians, and Asian Americans and Pacific Islanders.

No (1 point)

For more information, visit us at www.diabetes.org or call 1-800-DIABETES

Visit us on Facebook

Facebook.com/AmericanDiabetesAssociation

Add up your score.



Lower Your Risk

151:775-783, 2009.

The good news is that you can manage your risk for type 2 diabetes. Small steps make a big difference and can help you live a longer,

If you are at high risk, your first step is to see your doctor to see if additional testing is

Visit diabetes.org or call 1-800-DIABETES for information, tips on getting started, and ideas for simple, small steps you can take to help lower your risk.



CDC Prediabetes Screening Test



COULD YOU HAVE PREDIABETES?

Prediabetes means your blood glucose (sugar) is higher than normal, but not yet diabetes. Diabetes is a serious disease that can cause heart attack, stroke, blindness, kidney failure, or loss of feet or legs. Type 2 diabetes can be delayed or prevented in people with prediabetes through effective lifestyle programs. Take the first step. Find out your risk for prediabetes.

TAKE THE TEST—KNOW YOUR SCORE!

Answer these seven simple questions. For each "Yes" answer, add the number of points listed. All "No" answers are O points.

Are you a woman who has had a baby weighing more than 9 pounds at birth?

Do you have a sister or brother with diabetes?

Do you have a parent with diabetes?

Find your height on the chart. Do you weigh as much as or more than the weight listed for your height?

Are you younger than 65 years of age and get little or no exercise in a typical day?

Are you between 45 and 64 years of age?

Are you 65 years of age or older?

Add your score and check the back of this page to see what it means.

| P | II-KISK WEI | GHI CHAK | |
|--------|---------------|----------|---------------|
| Height | Weight Pounds | Height | Weight Pounds |
| 4'10" | 129 | 5'7" | 172 |
| 4'11" | 133 | 5'8" | 177 |
| 5'0" | 138 | 5'9" | 182 |
| 5'1" | 143 | 5'10" | 188 |
| 5'2" | 147 | 5'11" | 193 |
| 5'3" | 152 | 6'0" | 199 |
| 5'4" | 157 | 6'1" | 204 |
| 5'5" | 162 | 6'2" | 210 |
| 5'6" | 167 | 6'3" | 216 |
| | | 6'4" | 221 |

AT DICK MEICHT CHART

National Center for Chronic Disease Prevention and Health Promotion Division of Diabetes Translation



IF YOUR SCORE IS 3 TO 8 POINTS

This means your risk is probably low for having prediabetes now. Keep your risk low. If you're overweight, lose weight. Be active most days, and don't use tobacco. Eat low-fat meals with fruits, vegetables, and whole-grain foods. If you have high cholesterol or high blood pressure, talk to your health care provider about your risk for type 2 diabetes.

IF YOUR SCORE IS 9 OR MORE POINTS

This means your risk is high for having prediabetes now. Please make an appointment with your health care provider soon.

HOW CAN I GET TESTED FOR PREDIABETES?

Individual or group health insurance: See your health care provider. If you don't have a provider, ask your insurance company about providers who take your insurance. Deductibles and copays may apply.

Medicaid: See your health care provider. If you don't have a provider, contact a state Medicaid office or contact your local health department.

Medicare: See your health care provider. Medicare will pay the cost of testing if the provider has a reason for testing. If you don't have a provider, contact your local health department.

No insurance: Contact your local health department for more information about where you could be tested or call your local health clinic.





have prediabetes

You could be one of them.

Having prediabetes means you are at increased risk for developing serious health problems such as type 2 diabetes, stroke and heart disease.

You could have prediabetes if you have:

- High cholesterol **or**
- High blood pressure or
- A parent, brother or sister with diabetes

Your risk goes up if you are also overweight, and/or over age 45.

If you have prediabetes, we can help!

Ask your doctor how you can stop diabetes before it starts.







Are you at risk for prediabetes?



1 in 3 U.S. adults has prediabetes. Most don't know it. Are you at risk?

You may have prediabetes and be at risk for type 2 diabetes if you:

- · Are 45 years of age or older
- Are overweight
- Have a family history of type 2 diabetes
- Have high blood pressure
- Are physically active fewer than three times per week
- Ever had diabetes while pregnant (gestational diabetes) or gave birth to a baby that weighed more than 9 pounds

Prediabetes can lead to serious health problems

Having prediabetes means your blood glucose (sugar) level is higher than normal, but not high enough to be diagnosed as diabetes. But, nearly 90 percent of adults who have prediabetes don't know they have it.

If you have prediabetes and don't lose weight or increase your physical activity, you could develop type 2 diabetes within five years. Type 2 diabetes is a serious condition that can lead to health issues such as heart attack, stroke, blindness, kidney failure, or loss of toes, feet or legs.

What can you do?

Talk to your doctor about your risk of having prediabetes.

Here's the good news

If you have prediabetes, your doctor may refer you to a proven lifestyle change program that can help you prevent or delay getting type 2 diabetes.

The National Diabetes Prevention Program can help!

The National Diabetes Prevention Program (National DPP) uses a program that is proven to prevent or delay type 2 diabetes, and will help you lower your risk by improving your food choices and increasing physical activity.

How does it work? As part of a group in your community or online, you will work with a trained lifestyle coach to learn the skills you need to make lasting lifestyle changes. You will learn to eat healthy, add physical activity to your life, manage stress, stay motivated and solve problems that can get in the way of healthy changes.

Features

- Trained coach to guide and encourage you
- · In-person or online
- CDC-approved program
- Support from others working on the same goals as you
- Skills to help you lose weight, be more physically active and manage stress
- Some insurance companies will cover

What participants are saying ...

"I love having a lifestyle coach. She has given us great information, helped me stay on track and stay positive!"

—Bruce

"I'm so excited because I went to the doctor last week and all of my numbers were down and I officially no longer have prediabetes."

—Vivien

Now is the time to take charge of your health and make a change! Ask your doctor or nurse.







So you have prediabetes ... now what?



Prediabetes means your blood glucose (sugar) level is higher than normal, but not high enough to be diagnosed as diabetes. This condition raises your risk of type 2 diabetes, stroke and heart disease.

Some insurance plans will cover the cost of the program. Check with your insurance provider to see if it is covered. Also, some places that provide the program will adjust the fee you pay based on your income.

What can you do about it?

The good news is that there's a program that can help you.

The National Diabetes Prevention Program, led by the Centers for Disease Control and Prevention (CDC), uses a method proven to prevent or delay type 2 diabetes.

By improving food choices and increasing physical activity, your goal will be to lose 5 to 7 percent of your body weight—that is 10 to 14 pounds for a person weighing 200 pounds.

These lifestyle changes can cut your risk of developing type 2 diabetes by more than half.

How does the program work?

As part of a group, you will work with a trained diabetes prevention coach and other participants to learn the skills you need to make lasting lifestyle changes. You will learn to eat healthy, add physical activity to your life, manage stress, stay motivated and solve problems that can get in the way of healthy changes.

The program lasts one year, with 16 sessions taking place about once a week and six to eight more sessions meeting once a month. By going through the program with others who have prediabetes you can celebrate each other's successes and work together to overcome challenges.

Why should you act now?

Without weight loss and moderate physical activity, many people with prediabetes will develop type 2 diabetes within five years. Type 2 diabetes is a serious condition that can lead to health issues such as heart attack, stroke, blindness, kidney failure, or loss of toes, feet or legs. **NOW is the time to take charge of your health and make a change.**

Features of the program:

- · A trained coach to guide and encourage you
- A CDC-approved program
- Group support
- Skills to help you lose weight, be more physically active and manage stress

What participants are saying ...

"I love having a lifestyle coach. She has given us great information, helped me stay on track and stay positive!"

—Bruce

"I'm so excited because I went to the doctor last week and all of my numbers were down and I officially no longer have prediabetes."

-Vivien

Sign up today for a program near you!

To find a program in our area that is part of the National Diabetes Prevention Program, visit **cdc.gov/diabetes/ prevention.**

















Letter template

Use/adapt these templates to conduct efficient follow-up and referral with patients who have been identified as having prediabetes

<<YOUR LETTERHEAD>>

<<ADDRESS>>

<< PHONE NUMBER>>

<<DATE>>

<<PATIENT NAME>>

<< PATIENT ADDRESS>>

Dr. Mr./Mrs. << PATIENT LAST NAME>>,

Thank you for being a patient of the << PRACTICE NAME HERE>>. We are writing to tell you about a service to help make your health better.

Based on our review of your medical chart, you have a condition known as prediabetes. This means your blood sugar is higher than normal, which increases your risk of developing serious health problems including type 2 diabetes, as well as heart disease and stroke.

We have some good news. Our office wants you to know that you may be eligible for a diabetes prevention program run by our partners, << NAME OF PROGRAM PROVIDER>>. This program is proven to reduce your risk of developing diabetes and other health problems.

We have sent a referral to <<NAME OF PROGRAM PROVIDER>> and someone will call you to discuss the program, answer any questions you may have and, if you are interested, enroll you in the program.

Please feel free to give << NAME OF PROGRAM PROVIDER>> a call at << PHONE NUMBER>>.

-OR-

We have sent a referral to << NAME OF PROGRAM PROVIDER>> and we urge you to call << PHONE NUMBER>> to learn more about the program and enroll.

We hope you will take advantage of this program, which can help prevent you from developing serious health problems.

Sincerely,

Dr. << PHYSICIAN LAST NAME>>

















Sample "Talking points" for phone outreach

- Hello << PATIENT NAME>>.
- I am calling from << PRACTICE NAME HERE>>.
- I'm calling to tell you about a program we'd like you to consider, to help you prevent some serious health problems.
- Based on our review of your medical chart, you have a condition known as prediabetes. This means your blood sugar is higher than normal, which makes you more likely to develop serious health problems including type 2 diabetes, stroke and heart disease.
- We have some good news, too.
- You may be eligible for a diabetes prevention program run by our partners, << NAME OF PROGRAM PROVIDER>>.
 - Their program is based on research proven to reduce your risk of developing diabetes and other health problems.

Option A

- We have sent a referral to << NAME OF PROGRAM PROVIDER >> and someone will call you to discuss the program, answer any questions you may have and, if you are interested, enroll you in the program.
- Please feel free to give << NAME OF PROGRAM PROVIDER>> a call at << PHONE NUMBER>>.
- Do you have any questions for me?
- · Thank you for your time and be well.

Option B

- We have sent a referral to << NAME OF PROGRAM PROVIDER>> and we urge you to call << PHONE NUMBER>> to learn more about the program and enroll.
- We hope you will take advantage of this program, which can help prevent you from developing serious health problems.
- Do you have any questions for me?
- · Thank you for your time and be well.







Incorporate screening, testing and referral into practice





M.A.P. (Measure, Act, Partner)

patients with prediabetes and referring to community-based diabetes prevention programs. "Point-of-Care" and "Retrospective" methods may be used together or alone. THE M.A.P. (Measure, Act, Partner) to prevent type 2 diabetes—physicians and care teams can use this document to determine roles and responsibilities for identifying adult

Choose and check what works best for your practice

| Step 1: Measure | When | Who | How (draw from AMA-CDC tools) |
|---|--|--|--|
| Point-of-care method • Assess risk for prediabetes during routine office visit • Test and evaluate blood glucose level based on risk status | At the front deskDuring vital signs | ReceptionistMedical assistantNursePhysicianOther | Provide "Are you at risk for prediabetes?" patient education handout in waiting area Use/adapt "Patient flow process" tool Use CDC or ADA risk assessment questionnaire at check-in Display 8 x 11" patient-facing poster promoting prediabetes awareness to your patients Use/adapt "Point-of-care algorithm" |
| Retrospective method Query EHR to identify patients with BMI ≥24 and blood glucose level in the prediabetes range | • Every 6–12 months | Health IT staffOther | Use/adapt "Retrospective algorithm" |
| Step 2: Act | | | |
| Point-of-care method Counsel patient re: prediabetes and treatment options during office visit Refer patient to diabetes prevention program Share patient contact info with program provider* | During the visit | Medical assistantNursePhysicianOther | Advise patient using "So you have prediabetes now what?" handout Use/adapt "Health care practitioner referral form" Refer to "Commonly used CPT and ICD codes" |
| Retrospective method Inform patient of prediabetes status via mail, email or phone call Make patient aware of referral and info sharing with program provider Refer patient to diabetes prevention program Share patient contact info with program provider* | Contact patient soon after EHR query | Health IT staff Medical assistant (for phone calls) Other | Use/adapt "Patient letter/phone call" template Use/adapt "Health care practitioner referral form" for making individual referrals Use/adapt "Business Associate Agreement" template on AMA's website if needed |
| Step 3: Partner | | | |
| With diabetes prevention programs Engage and communicate with your local diabetes prevention program Establish process to receive feedback from program about your patients' participation | Establish contact before making 1st referral | Medical assistantNursePhysicianOther | Use/adapt " <u>Business Associate Agreement</u> " template on AMA's website if needed Refer to "Commonly used CPT and ICD codes" |
| With patients Explore motivating factors important to the patient At follow-up visit, order/review blood tests to determine impact of program and reinforce continued program participation Discuss program feedback with patient and integrate into care plan | During office visitOther | Office managerOther | Advise patient using "So you have prediabetes now what?" handout and provide CDC physical activity fact sheet www.cdc.gov/physicalactivity |

* To share patient contact information with a diabetes prevention program, you may need a Business Associate Agreement (BAA). PCMH recognition via Standard 4: Self-Care Support, B. Provide Referrals to Community Resources (3 points), NCQA Facilitating PCMH Recognition, 2011.) Following the M.A.P. for Preventing Type 2 Diabetes can help your practice achieve Patient Centered Medical Home (PCMH) recognition, as well as Meaningful Use of your electronic medical record. (Supports

The American Medical Association and the Centers for Disease Control and Prevention have created a tool kit that can help physician practices screen and refer patients to evidence-based diabetes prevention programs. Visit **preventdiabetesstat.org** to learn more. Physicians and other health care providers should also use their independent judgment when referring to a diabetes prevention program.

Sample patient flow process

MEASURE

CHECK-IN

- If age ≥18 and patient does not have diabetes, provide CDC Prediabetes Screening Test or ADA Diabetes Risk Test
- · Patient completes test and returns it
- Insert completed test in paper chart or note risk score in EMR

ROOMING/VITALS

- Calculate BMI (using table) and review diabetes risk score
- If elevated risk score or history of GDM, flag for possible referral

ACT

EXAM/CONSULT

- Follow "Point-of-care prediabetes identification algorithm"
- Determine if patient has prediabetes and BMI ≥24
 (≥ 22 for Asians) or a history of GDM
- Advise re: diet/exercise and determine willingness to participate in a diabetes prevention program
- If patient agrees to participate, proceed with referral

PARTNER

REFERRAL

• Complete and submit referral form via fax or email



FOLLOW UP

- Contact patient and troubleshoot issues with enrollment or participation
- † Use handouts included in this toolkit or request brochures from your local diabetes prevention program provider.







Point-of-care prediabetes identification

MEASURE

If patient is age ≥18 and does not have diabetes, provide self-screening test
(CDC Prediabetes Screening Test or ADA Diabetes Risk Test)

If self-screening test reveals risk, proceed to next step

Review medical record to determine if BMI ≥24 (≥22 if Asian) or history of GDM*



YES

If no: Patient does not currently meet program eligibility requirements

NO

Determine if a HbA1C, FPG or OGTT was performed in the past 12 month



Order one of the tests below:

- Hemoglobin A1C (HbA1C)
- Fasting plasma glucose (FPG)
- Oral glucose tolerance test (OGTT)





| Diagnostic test | Normal | Prediabetes | Diabetes |
|-------------------------------------|--------|-------------|----------|
| HbA1C(%) | < 5.7 | 5.7-6.4 | ≥ 6.5 |
| Fasting plasma glucose (mg/dL) | < 100 | 100–125 | ≥ 126 |
| Oral glucose tolerance test (mg/dL) | <140 | 140–199 | ≥ 200 |
| | | | |

ACT

Encourage patient to maintain a healthy lifestyle.

Continue with exam/ consult. Retest within three years of last negative test. Refer to diabetes prevention program, provide brochure.

Consider retesting annually to check for diabetes onset.

Confirm diagnosis; retest if necessary.

Counsel patient re: diagnosis.

Initiate therapy.

PARTNER

Communicate with your local diabetes prevention program.

Contact patient and troubleshoot issues with enrollment or participation. At the next visit, ask patient about progress and encourage continued participation in the program.

Adapted from: New York State Department of Health. New York State Diabetes Prevention Program (NYS DDP) prediabetes identification and intervention algorithm. New York: NY Department of Health; 2012.

*History of GDM = eligibility for diabetes prevention program







Referring patients to a diabetes prevention program

Method 1:

Point-of-care identification and referral

Download and display patient materials

Download and print the practice and patient resources included in this guide in advance of patient visits, so your office can have them available in the waiting room or during consult.

Measure

Step 1 – During check-in: If age ≥18 and patient does not have diabetes, give him/her the "<u>CDC Prediabetes Screening Test</u>" or American Diabetes Association "<u>Diabetes Risk Test</u>". After patient completes the test and returns it, insert completed test in the paper chart or note risk score in the electronic medical record (EMR). Screening test can also be mailed to patient along with other pre-visit materials.

Step 2 – During rooming/vitals: Calculate the patient's <u>body mass index</u>. Most EMRs can calculate BMI automatically. Review the patient's diabetes risk score and if elevated (\geq 5 on ADA test or \geq 9 on CDC test), flag for possible referral.

Step 3 – During exam/consult: Follow the "Point-of-care prediabetes identification algorithm" to determine if patient has prediabetes.

If the blood test results **do not** indicate prediabetes:

Encourage the patient to maintain healthy lifestyle choices. Continue with exam/consult.

Act

A. If the patient screens positive for prediabetes and has BMI <24 (<22 if Asian):

- Introduce the topic of prediabetes by briefly explaining what it is and its relation to diabetes (use the handout "So you have prediabetes ... now what?"). Review the patient's own risk factors.
- Emphasize the importance of prevention, including healthy eating, increased physical activity, and the elimination of risky drinking and tobacco use. (Visit the National Diabetes Education Program's GAME PLAN to Prevent Type 2 Diabetes for additional patient resources.)
- B. If the patient screens positive for prediabetes and has BMI ≥24 (≥22 if Asian):
 - Follow the steps in "A" above, discuss the value of participating in a diabetes prevention program, and determine the patient's willingness to let you refer him/her to a program.
 - If the patient agrees, complete and send the <u>referral form</u> to a community-based or online diabetes prevention program, depending on patient preference.
 - If patient declines, offer him/her a program handout and re-evaluate risk factors at next clinic visit.

Step 4 – Referral to diabetes prevention program: Most diabetes prevention programs are configured to receive referrals via conventional fax (over a phone line) or secure email. Complete the <u>referral form</u> and submit to a program as follows:

- A. If using a paper referral form, send via fax (over a phone line) or scan and email
- B. If the referral form is embedded in your EMR, either fax (over a phone line) or email using the EMR
 - Some diabetes prevention programs can also receive an e-fax (over the Internet)

Physicians and other health care providers should also use their independent judgment when referring to a diabetes prevention program.

Partner

Step 5 – Follow-up with patient: Contact patient and troubleshoot issues with enrollment or participation. At the next visit, ask patient about progress and encourage continued participation in the program.







Retrospective prediabetes identification

MEASURE

Query EMR or patient database every 6–12 months using the following criteria:

A. Inclusion criteria:

- Age ≥18 years and
- Most recent BMI ≥24 (≥22 if Asian) and
- A positive lab test result within previous 12 months:
 - HbA1C 5.7-6.4% (LOINC code 4548-4) or
 - FPG 100-125 mg/dL (LOINC code 1558-6) or
 - OGTT 140–199 mg/dL (LOINC code 62856-0) or
- History of gestational diabetes (ICD-9: V12.21)

B. Exclusion criteria:

- Current diagnosis of diabetes (ICD-9: 250.xx) or
- Current Insulin use

Generate a list of patient names with relevant information



ACT

Use the patient list to:

- A. Contact patients to inform of risk status, explain prediabetes, and share info on diabetes prevention programs, and/or
- B. Send patient info to diabetes prevention program provider
 - Program coordinator will contact patient directly, and
- C. Flag medical record for patient's next office visit



PARTNER

Discuss program participation at next visit







Retrospective identification and referral

Step 1 - Query EMR or patient database

Measure

Query your EMR or patient database every 6–12 months using the following criteria:

A. Inclusion criteria:

- Age ≥18 years and
- BMI ≥24 (≥22 if Asian) and
- A positive test result for prediabetes within the preceding 12 months:
 - HbA1C 5.7–6.4% or
 - Fasting plasma glucose 100–125 mg/dL or
 - Oral glucose tolerance test 140–199 mg/dL or
- Clinically diagnosed gestational diabetes during a previous pregnancy

B. Exclusion criteria:

- Current diagnosis of diabetes or
- Current Insulin use

Generate a list of patient names and other information required to make referrals:

Gender and birth dateMailing addressPhone number

Act

Step 2 – Referral to diabetes prevention program

- A. Contact patients via phone, email, <u>letter</u> or postcard to explain their prediabetes status and let them know about the diabetes prevention program.
- B. Send relevant patient information to your local (or online) diabetes prevention program coordinator and have him/her contact the patient directly (may require <u>Business Associate Agreement</u>).
- C. Flag patients' medical records for their next office visit.

Physicians and other health care providers should also use their independent judgment when referring to a diabetes prevention program.

Partner

During the next office visit, discuss diabetes prevention program participation:

- If the patient is participating, discuss program experience and encourage continued participation
- If the patient has declined to participate, stress the importance of lifestyle change and continue to encourage participation (use the handout "So you have prediabetes ... now what?")







Health care practitioner referral form to a diabetes prevention program

Send to: Fax: Email:

| PATIENT INFORMATION | | | | | | | | | | |
|---|--|--------------------------|--|--|--|--|--|--|--|--|
| First name | | | Address | | | | | | | |
| Last name | | | | | | | | | | |
| Health insurance | 1 | | City | | | | | | | |
| Gender □ N | ⁄lale □ Female | | State | | | | | | | |
| Birth date (mm/c | dd/yy) | | ZIP code | | | | | | | |
| Email | | | Phone | | | | | | | |
| By providing your information above, you authorize your health care practitioner to provide this information to a diabetes prevention program provider, who may in turn use this information to communicate with you regarding its diabetes prevention program. | | | | | | | | | | |
| PRACTITIONER INFO | DRMATION (COMPLETED BY H | IEALTH CARE PRACTITION | ER) | | | | | | | |
| Physician/NP/PA | | | Address | | | | | | | |
| Practice contact | | | City | | | | | | | |
| Phone | | | State | | | | | | | |
| Fax | | | ZIP code | | | | | | | |
| SCREENING INFORMATION | | | | | | | | | | |
| Body Mass Index (BMI) | | | | | | | | | | |
| Blood test (check | cone) | Eligible range | | Test result (one only) | | | | | | |
| ☐ Hemoglobin <i>A</i> | A1C | 5.7-6.4% | | | | | | | | |
| ☐ Fasting Plasma | a Glucose | 100-125 mg/dL | | | | | | | | |
| ☐ 2-hour plasma | a glucose (75 gm OGTT) | 140-199 mg/dL | | | | | | | | |
| Date of blood tes | st (mm/dd/yy): | | | | | | | | | |
| For Medicare req | uirements, I will maintair | n this signed original d | ocument in the patient's i | medical record. | | | | | | |
| Date | | Practitioner signatu | ıre | | | | | | | |
| 1 Y | By signing this form, I authorize my physician to disclose my diabetes screening results to the (insert program/organization name here) for the purpose of determining my eligibility for the diabetes prevention program and conducting other activities as permitted by law. | | | | | | | | | |
| | I understand that I am no authorization is voluntar | | oate in this diabetes scree | ning program and that this | | | | | | |
| 0 | • | | n at any time by notifying s taken before my physicia | my physician in writing. an received my written revocation. | | | | | | |
| | Date Pa | atient signature | | | | | | | | |

IMPORTANT WARNING: The documents accompanying this transmission contain confidential health information protected from unauthorized use or disclosure except as permitted by law. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless permitted to do so by law or regulation. If you are not the intended recipient and have received this information in error, please notify the sender immediately for the return or destruction of these documents. Rev. 05/30/14

BMI calculation chart

| Higher Hour Hour Hour Hour Hour Hour Hour Hou | | | | | | | | | | | | | | | | | | | | Ļ |
|---|--------|----------------|-----|-----|-----|------|-----|------|-----|------|------|-------|-------|-----|------|------|----|------|----|----------------------|
| 1 | 400 | 78 | 76 | 73 | 7 | 69 | 67 | 65 | 63 | 61 | 59 | 28 | 26 | 54 | 53 | 5 | 20 | 49 | 48 | greate |
| 1 1 1 1 2 2 2 2 2 2 | 390 | 76 | 74 | 72 | 69 | 29 | 65 | 63 | 61 | 29 | 28 | 26 | 22 | 53 | 52 | 20 | 49 | 48 | 46 | 1: 40 or |
| 1 1 1 1 2 2 2 2 2 2 | 380 | 74 | 72 | 70 | 29 | 9 | 63 | 62 | 09 | 28 | 26 | 22 | 53 | 52 | 20 | 49 | 48 | 46 | 45 | Obesity |
| 1 1 1 1 2 2 2 2 2 2 | 370 | 72 | 70 | 89 | 99 | 99 | 62 | 09 | 28 | 26 | 22 | 53 | 25 | 20 | 49 | 48 | 46 | 45 | 44 | xtreme |
| He control co | 360 | 11 | 89 | 99 | 99 | 62 | 09 | 28 | 27 | 22 | 23 | 25 | 20 | 49 | 48 | 46 | 45 | 44 | 43 | |
| He loo ii lo 120 ii lo 22 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | 350 | 69 | 99 | 64 | 62 | 09 | 28 | 22 | 22 | 23 | 25 | 20 | 49 | 48 | 46 | 45 | 44 | 43 | 42 | |
| HT 100 110 120 130 140 150 150 150 150 170 180 190 200 210 220 220 240 250 270 280 270 280 270 280 310 320 HT 1 12 21 22 24 26 28 30 32 32 34 36 37 39 41 42 45 46 46 85 50 51 53 55 57 59 61 HT 1 19 21 23 24 26 28 30 32 33 35 35 37 39 41 43 44 46 46 48 50 51 53 55 57 59 61 HT 1 19 21 23 24 25 24 25 28 29 31 33 35 37 39 41 43 44 46 46 48 50 51 53 55 57 59 61 HT 1 19 21 22 22 23 25 26 28 30 31 33 35 37 39 41 43 44 46 47 48 50 51 53 55 57 59 61 HT 1 19 21 22 23 24 25 24 25 27 29 31 33 35 37 39 40 42 44 46 47 49 51 53 55 57 59 51 HT 1 19 21 22 23 24 25 26 28 30 31 33 35 37 39 40 42 44 46 47 49 51 52 52 52 52 52 52 52 52 52 52 52 52 52 | 340 | 29 | 64 | 62 | 09 | 29 | 57 | 22 | 53 | 52 | 20 | 49 | 48 | 46 | 45 | 44 | 43 | 14 | 40 | |
| HT 100 110 120 120 120 120 120 140 150 160 170 180 190 200 210 220 240 250 260 270 260 270 200 200 200 200 200 200 200 200 20 | 330 | 99 | 63 | 19 | 29 | 22 | 22 | 53 | 25 | 20 | 49 | 47 | 46 | 45 | 44 | 42 | 41 | 40 | 39 | |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 250 270 290 290 290 290 290 290 290 290 290 29 | 320 | 63 | 19 | 26 | 22 | 22 | 53 | 25 | 20 | 46 | 47 | 46 | 45 | 44 | 42 | 41 | 40 | 39 | 38 | |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 260 270 280 290 290 290 290 290 290 290 290 290 29 | 310 | 61 | 29 | 22 | 23 | 53 | 52 | 20 | 46 | 47 | 46 | 45 | 43 | 42 | 14 | 40 | 39 | 38 | 37 | |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 220 240 250 250 250 270 280 HT 19 21 23 25 27 29 31 33 35 35 37 39 41 42 44 46 46 48 50 51 53 55 17 18 20 22 23 25 24 26 28 30 32 33 34 36 38 40 42 44 46 46 48 50 51 51 51 51 51 51 51 51 51 51 51 51 51 | 300 | 29 | 22 | 22 | 23 | 25 | 20 | 46 | 47 | 46 | 44 | 43 | 42 | 41 | 39 | 39 | 38 | 37 | 36 | Orange |
| 100 110 120 120 120 140 150 160 170 180 190 200 210 220 230 240 250 250 270 | 290 | 22 | 22 | 23 | 25 | 20 | 48 | 47 | 46 | 44 | 43 | 42 | 41 | 39 | 38 | 37 | 36 | 32 | 34 | |
| 100 110 120 130 140 150 150 170 180 190 200 210 220 240 250 260 | 280 | 55 | 53 | 51 | 20 | 48 | 47 | 45 | 44 | 43 | 41 | 40 | 39 | 38 | 37 | 36 | 32 | 34 | 33 | |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 260 260 260 260 270 270 270 270 270 270 270 270 270 27 | 270 | 53 | 21 | 20 | 48 | 46 | 45 | 44 | 42 | 41 | 40 | 39 | 38 | 37 | 36 | 32 | 34 | 33 | 32 | 6 |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 250 250 250 250 250 250 250 250 25 | 260 | 51 | 49 | 48 | 46 | 45 | 43 | 42 | 41 | 40 | 39 | 37 | 36 | 32 | 34 | 33 | 33 | 32 | 31 | - 29. |
| 19 | 250 | 49 | 47 | 46 | 44 | 43 | 45 | 40 | 39 | 38 | 37 | 36 | 32 | 34 | 33 | 32 | 31 | 31 | 30 | |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 220 11 | 240 | 47 | 45 | 44 | 43 | 14 | 40 | 39 | 38 | 37 | 36 | 32 | 34 | 33 | 32 | 31 | 30 | 59 | 53 | |
| HT 100 110 120 130 140 150 160 170 180 190 200 210 2 HT 19 21 23 25 27 29 31 33 35 37 39 41 HS 20 22 24 26 28 30 32 33 35 37 39 HT 19 21 23 25 24 26 28 30 31 32 34 36 HG 11 19 21 22 24 25 27 29 31 32 34 36 HG 11 19 21 22 24 25 27 29 31 32 34 36 HG 11 19 21 22 24 25 27 29 31 32 34 36 HG 11 19 21 22 24 25 27 29 31 32 34 36 HG 11 19 21 22 24 25 27 29 30 31 33 35 HG 11 19 21 22 24 25 27 29 30 31 33 35 HG 11 19 21 22 24 25 27 29 30 31 33 35 HG 12 14 16 17 18 20 22 23 25 26 28 29 31 HG 15 16 18 19 20 21 22 23 25 26 28 29 31 HG 15 16 18 19 20 21 22 23 25 26 28 29 31 HG 15 17 18 20 21 22 23 24 25 27 28 HG 15 17 18 19 20 21 22 23 25 26 28 29 HG 15 17 18 19 20 21 22 23 25 26 28 29 HG 17 19 20 21 22 23 25 26 28 29 HG 15 17 18 19 20 21 22 23 25 26 28 HG 17 19 20 21 22 23 25 26 28 HG 18 19 20 21 22 23 24 25 27 28 HG 19 10 11 23 24 25 27 28 HG 19 10 11 23 24 25 27 28 HG 19 20 21 22 23 24 25 27 28 HG 19 20 21 22 23 24 25 27 28 HG 19 20 21 22 23 24 25 27 28 HG 19 10 11 13 14 15 16 18 19 20 21 22 23 24 25 26 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 26 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 27 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 16 17 18 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 17 18 17 18 18 19 20 21 22 23 24 25 24 25 HG 11 13 14 15 17 18 17 18 18 19 20 21 22 23 24 25 24 | 230 | 45 | 44 | 42 | 41 | 40 | 38 | 37 | 36 | 32 | 34 | 33 | 32 | 31 | 30 | 30 | 59 | 28 | 27 | Yellov |
| HT 100 110 120 130 140 150 160 170 180 190 200 2 2 2 2 2 2 2 2 2 2 3 1 3 3 3 5 3 7 39 18 20 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | 220 | 43 | 42 | 40 | 39 | 38 | 37 | 36 | 35 | 34 | 33 | 32 | 3 | 30 | 53 | 58 | 28 | 27 | 56 | |
| 100 110 120 130 140 150 160 170 180 190 200 110 120 131 31 31 31 31 31 31 | 210 | 41 | 39 | 38 | 37 | 36 | 35 | 34 | 33 | 32 | 31 | 30 | 59 | 28 | 27 | 27 | 56 | 56 | 52 | |
| HT 100 110 120 130 140 150 160 17 19 21 23 25 27 29 31 18 20 22 24 26 28 30 17 19 21 23 24 26 28 30 17 19 21 22 24 25 27 16 18 20 21 23 24 25 27 16 18 20 21 23 24 25 27 16 18 20 21 22 24 25 27 16 18 20 21 22 24 25 27 16 17 19 20 21 22 24 17 19 20 21 22 24 18 19 21 22 24 19 14 15 17 19 20 21 11 13 14 15 16 18 19 21 12 14 15 16 18 19 20 12 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 | 200 | 39 | 37 | 36 | 32 | 34 | 33 | 32 | 31 | 30 | 59 | 28 | 28 | 27 | 56 | 25 | 24 | 24 | 24 | 5-24.9 |
| HT 100 110 120 130 140 150 160 17 19 21 23 25 27 29 31 18 20 22 24 26 28 30 17 19 21 23 24 26 28 30 17 19 21 22 24 25 27 16 18 20 21 23 24 25 27 16 18 20 21 23 24 25 27 16 18 20 21 22 24 25 27 16 18 20 21 22 24 25 27 16 17 19 20 21 22 24 17 19 20 21 22 24 18 19 21 22 24 19 14 15 17 19 20 21 11 13 14 15 16 18 19 21 12 14 15 16 18 19 20 12 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 | 190 | 37 | 36 | 34 | 33 | 32 | 31 | 30 | 59 | 28 | 28 | 27 | 56 | 22 | 22 | 24 | 23 | 23 | 22 | ght: 18.5 |
| HT 100 110 120 130 140 150 160 17 19 21 23 25 27 29 31 18 20 22 24 26 28 30 17 19 21 23 24 26 28 30 17 19 21 22 24 25 27 16 18 20 21 23 24 25 27 16 18 20 21 23 24 25 27 16 18 20 21 22 24 25 27 16 18 20 21 22 24 25 27 16 17 19 20 21 22 24 17 19 20 21 22 24 18 19 21 22 24 19 14 15 17 19 20 21 11 13 14 15 16 18 19 21 12 14 15 16 18 19 20 12 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 18 19 20 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 11 13 14 15 16 17 18 19 | 180 | 35 | 34 | 33 | 32 | 31 | 30 | 59 | 28 | 27 | 56 | 22 | 22 | 24 | 23 | 23 | 22 | 21 | 21 | thy Wei |
| HT 100 110 120 130 140 150 161 151 151 151 151 151 151 151 151 151 | 170 | 33 | 32 | 31 | 30 | 59 | 28 | 27 | 56 | 22 | 22 | 24 | 23 | 23 | 22 | 21 | 21 | 20 | 20 | |
| HT 100 110 120 130 140 111 11 11 11 12 12 12 12 12 12 12 12 12 | 160 | 31 | 30 | 59 | 28 | 27 | 56 | 22 | 22 | 54 | 23 | 23 | 22 | 21 | 21 | 20 | 19 | 19 | 19 | Gree |
| 11 100 110 120 130 130 131 131 131 131 131 131 131 13 | 150 | 56 | 28 | 27 | 56 | 25 | 25 | 24 | 23 | 22 | 22 | 77 | 77 | 20 | 19 | 19 | 18 | 18 | 17 | |
| HT 100 110 120 130 11 21 23 25 12 20 22 24 13 20 22 23 17 18 20 22 16 18 20 21 16 17 19 21 15 16 18 20 17 18 20 18 10 11 19 21 10 11 19 21 10 11 19 21 11 10 11 19 11 11 11 11 11 11 11 11 11 11 11 11 11 11 | 140 | 27 | 56 | 25 | 24 | 24 | 23 | 22 | 22 | 21 | 20 | 20 | 19 | 19 | 8 | 9 | 17 | 17 | 16 | 3.5 |
| 100 19 19 19 19 19 19 19 19 19 19 19 19 19 | 130 | 25 | 24 | 23 | 23 | 22 | 21 | 21 | 20 | 19 | 19 | 18 | 18 | 17 | 17 | 16 | 16 | 15 | 5 | |
| 100 19 19 19 19 19 19 19 19 19 19 19 19 19 | 120 | 23 | 22 | 22 | 21 | 20 | 20 | 19 | 18 | 18 | 17 | 17 | 16 | 16 | 15 | 15 | 14 | 14 | 14 | ght: Les |
| 100 19 19 19 19 19 19 19 19 19 19 19 19 19 | 110 | 21 | 20 | 20 | 19 | 18 | 18 | 17 | 17 | 16 | 16 | 15 | 15 | 14 | 14 | 14 | 13 | 13 | 13 | ıderwei _ş |
| 보 부 : . | 100 | 19 | 18 | 18 | 17 | 17 | 16 | 16 | 15 | 15 | 41 | 14 | 14 | 13 | 13 | 12 | 12 | 12 | = | |
| | WEIGHT | неіднт 5'0" | 5.1 | 2.5 | 5.3 | 5.4" | 2.2 | 2.6" | 2.7 | 2.8. | 2.6. | 2,10, | 5'11" | 0.9 | .1.9 | .2.9 | 9 | .4.9 | 9 | • |

BMI stands for "BODY MASS INDEX" which is an estimate of total body fat based on height and weight. It is used to screen for weight categories that may lead to health problems. THE GOAL for most people is to have a BMI in the green area. It is usually best for your BMI to stay the same over time or to gradually move toward the green area.

Codes: When screening for prediabetes and diabetes

| Codes for prediabetes and diabetes screening*.t.# | | | | | | | | |
|---|--------------------------------------|--|--|--|--|--|--|--|
| International Classification of Dise | ases (ICD)-9 for diabetes screening | Current Procedural Terminology (CPT®) for diabetes screening tests | | | | | | |
| V77.1 | Diabetes Screening | CPT 82947 | Fasting Plasma Glucose Test | | | | | |
| 790.2 | Abnormal Glucose | CPT 82950 | Post-meal Glucose (2-hour plasma glucose; 2hPG; 2 hr specimen) | | | | | |
| 790.21 | Impaired Fasting Glucose | CPT 82951 | Oral Glucose Tolerance (3 specimens with 2 hr value included) | | | | | |
| 790.22 | Impaired Glucose Tolerance (oral) | CPT 83036 | Hemoglobin A1C | | | | | |
| 790.29 | Other Abnormal Glucose NEC | CPT 83036QW | Hemoglobin A1C (used for POC test that is CLIA waived [~DCA]) | | | | | |
| 278.00 | Obesity | | | | | | | |
| 278.02 | Overweight | | | | | | | |

These codes may be useful to report services/tests performed to screen for prediabetes and diabetes.

References



^{*} New York State Department of Health. New York State Diabetes Prevention Program (NYS DPP) prediabetes identification and intervention algorithm. New York: NY Dept of Health; 2012.

[†] American Diabetes Association. Standards of medical care in diabetes—2013. Diabetes Care. January 2013; 36:S11-66. doi: 10.2337/dc13-S011

[‡] Ackermann RT. Coding Guide for Diabetes and Prediabetes Testing. 2013. (Published here with permission from Ronald T. Ackermann MD, MPH.)